



CULTIVATION SHEET

"Olive Tree"

Family: Oleaceae



New cultivations

PROVIDE FOR FERTIGATION DURING THE TRANSPLANTION STAGE

1:400 OR 250 ML/HL IN THE LAST IRRIGATION STAGE

FOLIAR TREATMENT	DOSE
POST PRUNING	1:300 OR 333 ML/HL
AFTER ADVERSE WEATHER EVENTS (FROSTS)	1:500 OR 200 ML/HL
VEGETATIVE DEVELOPMENT	
MONTHLY TREATMENT	1:500 OR 200 ML/HL
PLANTS WITH FERTIGATION SYSTEM	DOSE
MONTHLY TREATMENT	1:200 O 500 ML/HL

Repeat every 30 days distribute in the last irrigation phase

Ongoing cultivations

FOLIAR TREATMENT	DOSE
POST HARVEST	1:500 OR 200 ML/HL
POST PRUNING SPRAY ALSO ALL THE TRUNK	1:300 OR 333 ML/HL
VEGETATIVE DEVELOPMENT	1:500 OR 200 ML/HL
BETWEEN FRUIT SET AND VERAISON	1:500 OR 200 ML/HL
FRUIT ENLARGMENT	1:500 OR 200 ML/HL

Repeat every 30 days increase the frequency every 7 days to make the environment unwelcome to insects and parasites in the most critical periods

FERTIGATION	DOSE
DURING THE SPRING - SUMMER VEGETATIVE STAGES	1:250 OR 400 ML/HL
WURING THE AUTUMN - WINTER VEGETATIVE STAGES	1:300 OR 333 ML/HL

*Repeat every 15 - 20 days based on needs
Distribute in the last irrigation stage*



The Migliorini farm has chosen our Wood Distillate for its cultivation. Dr. Giuseppe Migliorini says that his olive trees are now free of olive fly and pathogenic attacks



<https://www.youtube.com/?v=Pa-HF8ITKW8>

Compatibility:

The product has no contraindications for miscibility with other formulations. It is advisable to carry out compatibility tests on crops on small surfaces before extending the treatment to larger areas.

Warnings:

To be used exclusively in agriculture. Whoever uses the product is responsible for any damage resulting from the improper use of the product. Compliance with the aforementioned instructions is an essential condition to ensure the effectiveness of the treatment and to avoid damage to plants, people, animals.

*For more information and customized protocols, please contact BioEsperia.